	Technical Data Sheet			
Use in	 Pharmaceutical Industry For industrial, laboratory & research applications only Basic medium according to EP 2.6.13 and USP <62> 			
Use for	 Examination of non-sterile products Test for specified micro-organisms Test for Salmonella spp. Application: Typically, 10 g of product are diluted in 90 mL Tryptic Soy Broth (TSB, e.g., artNo. 501.B090), mixed and incubated at 30-35 °C for 18-24 h. 0.1 mL is transferred to Rappaport-Vassiliadis Broth and incubated at 30-35 °C for 18-24 h. An aliquot is subcultured on XLD Agar and incubated at 30-35 °C for 18-48 h. Growth of reddish or red colonies, with or without black centre indicates the presence of Salmonella. In case such colonies are detected an identification test needs to prove the presence/absence of Salmonella. 			
Typical composition per liter	Xylose L-Lysine Na-deoxycholate Lactose x H₂O Sucrose Na-thiosulfate This medium can be adju performance criteria requ	7.5 g 7.5 g 6.8 g usted / or s	Ferric ammonium citrate Phenol red	3 g 5 g 0.8 g 80 mg 13.5 g
Irradiation	Not irradiated			
Filling volume	• 23-26 mL			
Packaging	 Single bagged, staples of 10 plates Transparent High barrier foil against desiccation 6 staples of 10 plates per packaging unit Temperature isolated handle-bag in the cardboard-boxes 			
Plates per box	60 plates (6 staples with 10 plates each)			
Shelf life	4 months from production date			



	Technical Data Sheet	
Storage conditions	 Recommended storage temperature: 15-25 °C Should be stored at temperatures as stable as possible Store protected from light exposure Before use: it is recommended to keep the plates upright (agar on the lower part, lid on the upper part) to avoid formation of extra condensation After use: it is recommended to keep the plates upside down (agar on the upper part, lid on the lower part) to reduce the risk of accumulation of condensation during incubation which can affect colony formation 	
Label	On the side, at the bottom	
Label information	 Product name: XLD Expiry date: YYYYMMMDD → MMM in letters (e.g.: 2023Nov04) Lot-number Individual number Barcode 	
Barcode	 2-dimensional (data matrix), 20 digits: Digits 1-3: ArtNo. Digits 4-9: Lot-Number Digits 10-14: Individual-Number Digits 15-20: Date (YYMMDD) 	
Delivery	 Temperature controlled delivery on request For shipments of larger amounts plastic pallets in Euro-size can be used 	
Petri dish	 Locking lid 90 mm plate, made from polystyrene Long incubations possible – due to high filling volume Long expositions possible – due to specific design of plate Incubations in vent and closed position possible 	
Lid positions	 All plates are delivered in the non-locked position The plate contains 2 locked positions. If turning the lid clockwise the locked positions are in the following order: 1. Vent position 2. Closed position 	
Aerobic incubation	 The closed position provides ideal incubation conditions for aerobic microorganisms and limits the dehydration of the agar during incubation For long incubation of aerobic microorganisms, the closed position is recommended To lock the lid in the closed position turn the lid clockwise into the final stop position 	
Production	Production of selective media is made at the beginning of every quarter. To get the longest shelf life possible, we do recommend to place orders for delivery at this period of time.	



	Technical Data Sheet
Place of production	PharmaMedia Dr. Müller GmbH Gustav-Throm-Str. 1, 69181 Leimen - Germany

	Quality control, Certificates					
	Each lot of product can be obtained with a certificate of analysis (CoA):					
	Physico-chemical test parameters:				1	
	Appearance Clear, red, possibly with white crystals					
	pH value	7.2 – 7.6	-	-		
	Filling volume	23 – 26 mL				
	Growth Promotion test: 10-100 CFU*					
	S.Typhimurium	ATCC 14028	30-35 °C	16-18 h	50-200%	
	S.Abony	NCTC 6017	30-35 °C	16-18 h	50-200%	
	Inhibition test:	10,000-100,000	CFU			
	S.aureus	ATCC 6538	30-35 °C	48-52 h	No growth	
Certificates						
oci tilloutes	Indicative prop			40.041	1	_
	S.Typhimurium		30-35 °C	18-24 h	1	
	Colorless to pale	e pink colonies, c	colonies mos	ily with blac	ck center,	
	S.Abony	NCTC 6017	30-35 °C	18-24 h		_
	Colorless to pale				ck center.	
	medium red					
					T	
	Sterility control				Conform	
	recovery rate ag qualitative comp However, during in reference to a	parison with a programmer of a previously release	ctive referen evious appro at PMM reco ased batch o	ce) is requived batch is very rate date of the production of the production is required.	red – only the s requested. ata are determ ot.	nined
BSE policy	transmitting ar medicinal prod specified anima We neither sto infectivity tissu	with the current nimal spongiform lucts, we check al source, the coore or process rules (IA) nor rum a countries or reg	encephalop the CoO of ountry of origuminant raw inant raw inant raw ma	eathy via huraw materia in and the i materials o aterials who	uman or veter al in respect t nfectivity cate obtained from ose animal so	rinary to the egory. high ource



	Quality control, Certificates
Temperature stress	 Art. 470.0060 has been exposed to temperature stress conditions (3 days at 2-8 °C as well as 3 days at 30-35 °C) and has passed shelf-life testing at least 30 days after the assigned expiry date. Shelf-life testing comprise all regular tests which are part of the normal release test of this article except for sterility control (see CoA).

	Safety Data	
Toxic ingredients	• None	
Basic composition	See typical composition	
Solvent content	• None	
Safety data sheet required	Not mandatorily required	