

*Burkholderia cepacia* Complex Selective Agar 90 mm CS

Technical Data Sheet													
<b>Use in</b>	<ul style="list-style-type: none"> <li>Pharmaceutical Industry</li> <li>For industrial, laboratory &amp; research applications only</li> <li>Basic medium according to USP &lt;60&gt;</li> </ul>												
<b>Use for</b>	<ul style="list-style-type: none"> <li>Examination of non-sterile products</li> <li>Test for specified micro-organisms</li> <li>Test for <i>Burkholderia cepacia</i> complex</li> </ul> <p><b>Application:</b> Prepare a sample using a 1-in-10 dilution of not less than 1 g of the product to be examined. Use 10 mL or the quantity corresponding to 1 g or 1 mL to inoculate a suitable amount (determined as described in Suitability of the Test Method) of Soybean–Casein Digest Broth (e.g., art-No. 500.B100) or an appropriate dilution of Soybean–Casein Digest Broth as determined during method suitability (for example, a 1:10 dilution may be required when conducting optional testing of pharmaceutical waters). Then mix and incubate at 30-35 °C for 48-72 h.</p> <p>Subculture by streaking on a plate of <b><i>Burkholderia cepacia</i> Complex Selective Agar</b> and incubate at 30-35 °C for 48-72 h.</p> <p>Growth of colonies indicates the presence of <i>Burkholderia cepacia</i> complex. Any growth on BCCSA is confirmed by identification tests.</p>												
<b>Typical composition per liter</b>	<table> <tbody> <tr> <td>Caseine peptones</td> <td>10 g</td> <td>Yeast extract</td> <td>1.5 g</td> </tr> <tr> <td>Sucrose</td> <td>10 g</td> <td>Agar</td> <td>14 g</td> </tr> <tr> <td>NaCl</td> <td>5 g</td> <td>Selective supplements</td> <td></td> </tr> </tbody> </table> <p>This medium can be adjusted / or supplemented according to the performance criteria required.</p>	Caseine peptones	10 g	Yeast extract	1.5 g	Sucrose	10 g	Agar	14 g	NaCl	5 g	Selective supplements	
Caseine peptones	10 g	Yeast extract	1.5 g										
Sucrose	10 g	Agar	14 g										
NaCl	5 g	Selective supplements											
<b>BCCSA composition</b>	<p>BCCSA composition has been modified compared to the one recommended in the USP to ensure superior performances throughout the shelf life, a long shelf life and to keep inhibiting properties stable. The basic recipe described in USP &lt;60&gt; is used; however, following modifications have been introduced:</p> <ul style="list-style-type: none"> <li>Removal of <b>lactose</b>: <i>Burkholderia</i> sp. does not metabolize lactose. This sugar only supports the growth of the background flora</li> <li>Removal of <b>phenol red</b>: colonies growing on BCCSA need to be identified in any case, which makes the use of a colour indicator not relevant for the detection. Moreover, this red colour may not be stable with time</li> <li>Removal of <b>crystal violet</b>: crystal violet is inhibiting Gram + bacteria – however, as the antibiotic mix is inhibiting Gram + bacteria reliably, crystal violet can be omitted</li> <li>Antibiotics mix: optimized by PMM to ensure the inhibition of most of the background flora without inhibiting <i>Burkholderia</i> sp., which could be the case with gentamicin, vancomycin and polymyxin B</li> </ul>												

<b>Technical Data Sheet</b>	
<b>Irradiation</b>	<ul style="list-style-type: none"> <li>• Not irradiated</li> </ul>
<b>Filling volume</b>	<ul style="list-style-type: none"> <li>• 28-32 mL</li> </ul>
<b>Packaging</b>	<ul style="list-style-type: none"> <li>• Single bagged, staples of 10 plates</li> <li>• Transparent</li> <li>• High barrier foil against desiccation</li> <li>• 6 staples of 10 plates per packaging unit</li> <li>• Temperature isolated handle-bag in the cardboard-boxes</li> </ul>
<b>Plates per box</b>	<ul style="list-style-type: none"> <li>• 60 plates (6 staples with 10 plates each)</li> </ul>
<b>Shelf life</b>	<ul style="list-style-type: none"> <li>• 6 months from production date</li> </ul>
<b>Storage conditions</b>	<ul style="list-style-type: none"> <li>• Recommended storage temperature: 15-25 °C</li> <li>• Should be stored at temperatures as stable as possible</li> <li>• Store protected from light exposure</li> <li>• Before use: it is recommended to keep the plates upright (agar on the lower part, lid on the upper part) to avoid formation of extra condensation</li> <li>• After use: it is recommended to keep the plates upside down (agar on the upper part, lid on the lower part) to reduce the risk of accumulation of condensation during incubation which can affect colony formation</li> </ul>
<b>Label</b>	<ul style="list-style-type: none"> <li>• On the side, at the bottom</li> </ul>
<b>Label information</b>	<ul style="list-style-type: none"> <li>• Product name: BCCSA</li> <li>• Expiry date: YYYYMMDD → MMM in letters (e.g.: 2026Nov04)</li> <li>• Lot-number</li> <li>• Individual number</li> <li>• Barcode</li> </ul>
<b>Barcode</b>	<ul style="list-style-type: none"> <li>• 2-dimensional (data matrix), 20 digits:</li> <li>• Digits 1-3: Art.-No.</li> <li>• Digits 4-9: Lot-Number</li> <li>• Digits 10-14: Individual-Number</li> <li>• Digits 15-20: Date (YYMMDD)</li> </ul>
<b>Delivery</b>	<ul style="list-style-type: none"> <li>• Temperature controlled delivery on request</li> <li>• For shipments of larger amounts plastic pallets in Euro-size can be used</li> </ul>
<b>Petri dish</b>	<ul style="list-style-type: none"> <li>• Locking lid 90 mm plate, made from polystyrene</li> <li>• Long incubations possible – due to high filling volume</li> <li>• Long expositions possible – due to specific design of plate</li> <li>• Incubations in vent and closed position possible</li> </ul>

Technical Data Sheet	
<b>Lid positions</b>	<ul style="list-style-type: none"> <li>All plates are delivered in the non-locked position</li> <li>The plate contains 2 locked positions. If turning the lid clockwise the locked positions are in the following order:               <ol style="list-style-type: none"> <li>Vent position</li> <li>Closed position</li> </ol> </li> </ul>
<b>Aerobic incubation</b>	<ul style="list-style-type: none"> <li>The closed position provides ideal incubation conditions for aerobic microorganisms and limits the dehydration of the agar during incubation</li> <li>For long incubation of aerobic microorganisms, the closed position is recommended</li> <li>To lock the lid in the closed position, turn the lid clockwise into the final stop position</li> </ul>
<b>Production</b>	Production of selective media is made at the beginning of every quarter. To get the longest shelf life possible, we do recommend placing orders for <u>delivery</u> during this period.
<b>Place of production</b>	PharmaMedia Dr. Müller GmbH Gustav-Throm-Str. 1, 69181 Leimen - Germany

Quality control, Certificates		
<b>Certificates</b>	Each lot of product can be obtained with a certificate of analysis (CoA):	
	<b>Physico-chemical test parameters:</b>	
	Appearance	Clear, slightly yellowish
	pH value	6.6 – 7.0
	Filling volume	28 – 32 mL
	<b>Growth Promotion test: 10-100 CFU*</b>	
	<i>B. cepacia</i>	ATCC 25416    30-35 °C    44-48 h    50-200%
	<i>B. cenocepacia</i>	ATCC BAA-245    30-35 °C    48-72 h    50-200%
	<i>B. multivorans</i>	NCTC 13007    30-35 °C    44-48 h    50-200%
	<b>Inhibition test: 100-1,000 CFU</b>	
	<i>S. aureus</i>	ATCC 6538    30-35 °C    72-76 h    No growth
	<i>P. paraeruginosa</i>	ATCC 9027    30-35 °C    72-76 h    No growth
<b>Sterility control</b>	Conform	
<p>*According to Ph. Eur. 2.6.13 and USP&lt;62&gt;, no quantitative test (determination of recovery rate against a non-selective reference) is required – only the qualitative comparison with a previous approved batch is requested. However, during quality control at PMM recovery rate data are determined in reference to a previously released batch of the product.</p>		

<b>Quality control, Certificates</b>	
<b>Certificate of origin</b>	<p>All media lots produced by PMM can be obtained with a Certificate of Origin (CoO). All animal derived raw materials are specified as follows:</p> <ul style="list-style-type: none"> <li>• Raw material</li> <li>• Tissue</li> <li>• Animal source</li> <li>• Country of origin</li> <li>• Infectivity category (acc. to TSE guideline: EMA/410/01 current version)</li> </ul>
<b>BSE policy</b>	<ul style="list-style-type: none"> <li>• In compliance with the current note for guidance on minimizing the risk of transmitting animal spongiform encephalopathy via human or veterinary medicinal products, we check the CoO of raw material in respect to the specified animal source, the country of origin and the infectivity category. We neither store or process ruminant raw materials obtained from high infectivity tissues (IA) nor ruminant raw materials whose animal source originates from countries or regions with an undetermined risk (cat C/GBR IV).</li> </ul>

<b>Safety Data</b>	
<b>Toxic ingredients</b>	<ul style="list-style-type: none"> <li>• None</li> </ul>
<b>Basic composition</b>	<ul style="list-style-type: none"> <li>• See typical composition</li> </ul>
<b>Solvent content</b>	<ul style="list-style-type: none"> <li>• None</li> </ul>
<b>Safety data sheet required</b>	<ul style="list-style-type: none"> <li>• Not mandatorily required</li> </ul>